

Document Header Id #: 47473

RECEIVED IN PAD

Project #: G-35-W35

Cost share #:

Center #: 10/24-6-R0498-0A0

Center shr #:

OCA file #:

Project type: RES

Contract #: N00014-97-1-0879

Award type: GRANT

Prime #:

Contract entity: GTRC

CFDA: 12.300

PE #: N/A

Project unit: E & A SCI

Unit code: 33

Project director(s):

PDPI- VAN CAPPELLEN

P S

E & A SCI

(404)894-3883

Sponsor : NAVY/OFC OF NAVAL RESEARCH

Division Id: 103

/ 3314

Award period: 01-JUN-1997 to 30-SEP-1997 (performance) 30-SEP-1997 (reports)

Sponsor amount	New this change	Total to date
Contract value:	33,956.00	33,956.00
Funded:	33,956.00	33,956.00
Cost sharing amount:	0.00	0.00

Does subcontracting plan apply? N

Title: QUANTITATIVE CHEMICAL MASS TRANSFER IN COASTAL SEDIMENTS DURING...

PROJECT ADMINISTRATIVE DATA

OCA contact: Jacquelyn L. Bendall

(404) 894-4820

Sponsor technical contact:

DR. JOSEPH KRAVITZ

OFFICE OF NAVAL RESEARCH

MARINE AND GEOLOGY AND GEOPHYSICS

800 NORTH QUINCY STREET

ARLINGTON

VA

22217-5660

Phone: 703/696-4070

Fax:

Email:

Sponsor issuing office:

DIANE GALES

OFFICE OF NAVAL RESEARCH

ONR 252, BALLSTON CENTRE TOWER ONE

800 NORTH QUINCY STREET

ARLINGTON

VA

22217-5660

Phone: 703/696-2595

Fax:

Email:

Security class (U,C,S,TS): U

Defense priority rating : N/A

ONR resident rep is ACO (Y/N): Y

Supplemental sheet: ONR

Equipment title vests with: GIT

Administrative comments -

Initiation of grant.

u
(1)

Closeout Notice Date 29-OCT-1997

Project Number G-35-W35

Doch Id 47473

Center Number 10/24-6-R0498-0A0

Project Director VAN CAPPELLEN, PHILIPPE

Project Unit E & A SCI

Sponsor NAVY/OFC OF NAVAL RESEARCH

Division Id 3314

Contract Number N00014-97-1-0879

Contract Entity GTRC

Prime Contract Number

Title QUANTITATIVE CHEMICAL MASS TRANSFER IN COASTAL SEDIMENTS
DURING...

Effective Completion Date 30-SEP-1997 (Performance) 30-SEP-1997 (Reports)

Closeout Action:	Y/N	Date Submitted
Final Invoice or Copy of Final Invoice	Y	
Final Report of Inventions and/or Subcontracts	Y	
Government Property Inventory and Related Certificate	Y	
Classified Material Certificate	N	
Release and Assignment	N	
Other	N	

Comments

Distribution Required:

Project Director/Principal Investigator	Y
Research Administrative Network	Y
Accounting	Y
Research Security Department	N
Reports Coordinator	Y
Research Property Team	Y
Supply Services Department	Y
Georgia Tech Research Corporation	Y
Project File	Y

NOTE: Final Patent Questionnaire sent to PDPI

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 22 October 1997	3. REPORT TYPE AND DATES COVERED FINAL REPORT 01-JUN-97 to 30-SEP-97		
4. TITLE AND SUBTITLE Quantitative Chemical Mass Transfer in Coastal Sediments during Early Diagenesis			6. FUNDING NUMBERS N00014-97-1-0879	
6. AUTHOR(S) Philippe Van Cappellen				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Earth & Atmospheric Sciences Georgia Institute of Technology Atlanta, GA 30332-0340			8. PERFORMING ORGANIZATION REPORT NUMBER	
8. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Ballston Centre Tower One 800 North Quincy Street Arlington, VA 22217-5660			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
9. DISTRIBUTION / AVAILABILITY STATEMENT			12. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) This report summarizes activities related to ONR grant N00014-97-1-0879 to the Georgia Tech Research Corporation, for the period 1 June 1997 through September 30 1997 (Phase I). The work during Phase I has focused on (1) a sensitivity analysis of the effects of irrigation on the biogeochemistry of coastal marine sediments, 2) the upgrading of the existing multicomponent reactive transport model for early diagenesis, and (3) the preliminary development of a stochastic algorithm to represent burrowing activities of benthic macrofauna.				
14. SUBJECT TERMS Early diagenesis; sediments; biogeochemistry; bio-irrigation; reactive transport modeling			15. NUMBER OF PAGES 6 (+title page)	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

Office of Naval Research

Grant N00014-97-1-0879

**Quantitative Chemical Mass Transfer in Coastal
Sediments during Early Diagenesis**

Effects of Biological Transport, Mineralogy, and Fabric

FINAL REPORT

Principal Investigator

Philippe Van Cappellen

School of Earth and Atmospheric Sciences
Georgia Institute of Technology
Atlanta, GA 30332-0340
tel. (404) 894 3883
pvc@aqchem.eas.gatech.edu

Period: 1 June 1997 through 30 September 1997

INTRODUCTION

This report summarizes activities related to ONR grant N00014-97-1-0879 to the Georgia Tech Research Corporation, for the period 1 June 1997 through September 30 1997 (Phase I). The work during Phase I has focused on (1) a sensitivity analysis of the effects of irrigation on the biogeochemistry of coastal marine sediments, 2) the upgrading of the existing multicomponent reactive transport model for early diagenesis, and (3) the preliminary development of a stochastic algorithm to represent burrowing activities of benthic macrofauna.

ACHIEVEMENTS

Reactive transport code

The basis for the modeling efforts is STEADYSED1, a multi-component reactive transport code for steady-state early diagenesis which fully incorporates the reaction couplings among the reactive species of C, O, N, S, Fe and Mn. The model has been tested against extensive data sets collected by Canfield *et al.* (1993a, b) at three coastal marine sites that exhibit high rates of combined iron and manganese (hydr)oxide reduction. Results show that the model provides a consistent explanation of the entire body of multi-component, multi-site observations. The measured concentration profiles of 28 individual pore water and solid sediment species are satisfactorily reproduced. Furthermore, the model predicts the observed distributions of sulfate reduction rates, as well as diagnostic features of the pore water pH profiles.

The coupled nature of the reactive species distributions constrains the values of reaction and transport parameters, because a single set of parameters must account for the distributions of all the species at a given site. The parameters are further separated into site-dependent (e.g., deposition fluxes, bottom water composition, mixing coefficient, porosity) and reaction-specific parameters (e.g., rate coefficients, apparent equilibrium constants, limiting substrate concentrations). When fitting model-calculated distributions to observed ones, care is taken to minimize variations of the reaction-specific parameters from one site to another. This modeling strategy emphasizes extraction of reaction parameters from field data that are mechanistically meaningful.

Effects of irrigation


A sensitivity analysis was performed with STEADYSED1, under boundary conditions representative of coastal marine depositional environments. Conditions were chosen on the basis of previous work (Van Cappellen *et al.*, 1993; Tromp *et al.*, 1995; Middelburg *et al.*, 1997). The results show that model-calculated benthic exchange fluxes of most solutes are dominated by irrigation. Because of high rates of molecular oxygen reduction in organic-rich coastal sediments, the contribution of molecular diffusion to benthic uptake of O₂ may in some cases be of the same order of magnitude or even larger than that of irrigation. For nitrate,

molecular diffusion and irrigation cause exchange fluxes in opposite directions. As a consequence, there is a net transfer of nitrate from the water column to the sediments, although the interfacial pore water gradients predict diffusional fluxes out of the sediments. The results highlight the fundamental importance of pore water irrigation by macrofauna in controlling solute exchanges between sediments and water column in coastal environments.

Benthic fluxes and chemical conditions in sediments not only depend on the integrated intensity of irrigation, but also on the depth distribution of the irrigation exchange coefficient. This is shown in Figure 1, where three different distributions of the vertical irrigation exchange coefficient, α (in units of yr^{-1}), are considered. In two cases (1 & 3), the irrigation coefficient is assumed to decay exponentially with depth. This is the distribution used, for example, by Martin and Banta (1992) to reproduce tracer distributions of pore water bromide and $^{222}\text{Rn}/^{226}\text{Ra}$ disequilibrium in Buzzards Bay, Massachusetts. In the other case (2), the depth distribution of α follows a step function, that is, a uniformly irrigated surface layer overlies non-irrigated sediment. The depth integral of α is the same for all three distributions. Thus, the simulations compare sediments experiencing the same overall irrigation intensity.

The calculated pH profiles show distinct differences between the three irrigation scenarios (Figure 1). (Note: pH is one of the most sensitive parameters of sediment biogeochemistry.) The differences are most pronounced when comparing the step function to the exponential distributions. In the depth range 2-3 cm, the predicted pH values may deviate from one another by as much as one pH unit, i. e. the equivalent of an order of magnitude in hydrogen ion concentration.

Figure 1. Irrigation scenarios: (a) imposed depth distributions of the irrigation exchange coefficient; (b) calculated pH distributions. Calculations were performed under conditions typical of a coastal marine sediment. Boundary conditions and parameter values are the same in the three simulations, except for the distribution of the irrigation exchange coefficient. See text for further discussion.



Existing early diagenetic models use idealized depth distributions of the irrigation coefficient such as those presented in Figure 1. Nonetheless, little data or theoretical arguments have been presented for selecting one distribution over another (for exceptions, see: Martin and Banta, 1992; Wang and Van Cappellen, 1996). The sensitivity analysis performed here, however, shows that the depth distribution of the exchange coefficient has a significant impact on sediment biogeochemistry. A large part of the work planned for Phase II of the project will be devoted to the development of better theoretical and experimental constraints on the functional dependence of α on depth below the water-sediment interface.

CaCO₃ and pH buffering

Dissolution of deposited biogenic CaCO₃ or precipitation of authigenic CaCO₃ may exert a critical control on pore water pH, alkalinity and total dissolved inorganic carbon (e.g., Gaillard et al., 1989; Cai et al., 1995). The published version of STEADYSED1 (version 1.1) does not include the dissolution and precipitation reactions of CaCO₃. In the absence of buffering by these reactions, the model simulations tend to overpredict the magnitude of pore water pH fluctuations.

STEADYSED1 was upgraded (version 1.2) by including the rates of CaCO₃ dissolution and precipitation in the continuity equations of Ca²⁺, CaCO₃, total inorganic carbon (TC) and total alkalinity (Alk). As before, the pH profile is calculated by assuming that the dissolved carbonate system remains in local equilibrium. This is justified by the fast homogeneous acid-base kinetics. Several rate models for CaCO₃ are available for the user to choose from, as summarized in Table 1. Preliminary results show that by explicitly including CaCO₃ dissolution, the pore waters are buffered against large decreases in pH. More simulations are needed to fully assess the effects of CaCO₃ precipitation.

Table: Options for CaCO₃ dissolution/precipitation kinetics in Steadysed1 (v. 1.2)

Model	Rate expression	Reference
Empirical	$R_d = k_d[CaCO_3](1 - \Omega)^{4.5}$ $R_p = k_p(\Omega - 1)^{2.2}$	R_d : Cai <i>et al.</i> (1995) R_p : Zhong and Mucci (1993)
Mechanistic I	$R_d = k_1 a_{H^+} + k_2 a_{H_2CO_3} + k_3$ $R_p = k_{-1} a_{Ca^{2+}} a_{HCO_3^-} + k_{-2} a_{Ca^{2+}} a_{HCO_3^-}^2$ $+ k_{-3} a_{Ca^{2+}} a_{HCO_3^-}$	Chou <i>et al.</i> (1989)
Mechanistic II	$R_{total} = k_1[H^+] + k_2[H_2CO_3^0] + k_3[H_2O]$ $- k_4[Ca^{2+}][HCO_3^-]$	Plummer <i>et al.</i> (1979)
Mechanistic III (surface reaction)	$R_{total} = k_1 > CO_3^- a_{H^+}^2 + (k_2 - k_5) > Ca^+ a_{H_2CO_3}$ $+ k_4 - (k_6 - k_3) > CO_3 H^0 a_{CaHCO_3^+}$ $- k_7 > Ca^+ a_{H_2CO_3} a_{CaCO_3^0} - k_8 a_{CaCO_3^0}$	Arakaki and Mucci (1995)

R_d : dissolution rate; R_p : precipitation rate; R_{total} : overall reaction rate; a_i : activity of solute i ; $[i]$: concentration of solute i ; k_i : rate constant; $>i$: density of surface species i

The new version of the reactive transport code has not yet been made publicly available because some numerical problems remain to be solved. These problems are the consequence of the fast rates of CaCO_3 dissolution and precipitation, relative to the other kinetic processes represented in the model (e.g., the degradation of organic matter). Large differences among rate terms in the continuity equations are prone to cause numerical instabilities in the finite difference solutions. We are currently implementing changes in the code that counter these instabilities.

Stochastic burrowing

First steps have been taken in the development of a two-dimensional (2-D) stochastic model for random vertical burrowing. In this model the water-sediment interface is represented by a set of equally spaced spatial nodes. The algorithm considers each node in succession and decides whether or not to initiate burrowing. The decision is based on a randomly selected number, whereby the relative range of values leading to burrowing is determined by the average normalized burrow density.

The burrowing strategy plus environmental constraints that direct the activity of the macrofauna are translated into a probability distribution for the burrow length. Once a node has been selected as the site of a burrow, the length is determined by generating another random number. The value ranges assigned to different lengths are proportional to the corresponding probabilities. Based on numerous realizations of the stochastic burrow model, we calculate the average depth distribution of available burrow surface area (a proxy for the exchange coefficient), plus the deviation from the average. In this manner a statistical, rather than a deterministic, description of sediment burrowing is achieved. Our efforts in the near future will focus on the formulation of realistic probability distributions for burrow depths.

OTHER ACTIVITIES

A meeting of the P.I.s and the Program Officer was held at the Naval Research Laboratory in Stennis on September 26, 1997. Progress to date by the different laboratories was reviewed, and plans for Phase II (12/1/97 to 9/30/99) were discussed. Emphasis was placed on strengthening the integration of the P.I.s' efforts.

Work has begun on a manuscript on the effects of bio-irrigation on solute exchanges using data from the June 97 cruise to the Dry Tortugas organized by NRL (Lavoie, chief scientist). Dr. Furukawa is the lead author of this collaborative effort.

An abstract based on research closely related to that supported by this grant was submitted to the special session on *Diagenesis in Coastal and Marginal Sediments* of the 1998 Ocean Sciences Meeting (San Diego, February 9-13, 1998). It is entitled "*Spatial and Temporal Variability of the Iron and Sulfur Cycles in Salt*

Marsh Sediments: An Integrated Geochemical and Microbiological Study", by P. Van Cappellen, A. Roychoudhury, E. Viollier, K. Lowe, T. DiChristina, T. and J. Kostka.

The following activities are in the planning stage; they will enhance the visibility of research activities supported by ONR grant N00014-97-1-0879.

- A series of invited lectures by the P. I. at the Graduate College of GEOMAR (Forschungszentrum für marine Geowissenschaften) in Kiel, Germany (Nov. 26-28, 1997). The lectures will focus on the theoretical, numerical and experimental aspects of reactive transport in marine sediments.
- The organization by the P. I. of a thematic session entitled "*Biogeochemical Dynamics: From Molecular Approaches to Predictive Models*" at the Conference on Southeast Coastal Ocean Research (SECOR) (Savannah, April 7-10, 1998).

PERSONNEL

In addition to the P. I., first-year graduate student Chao Liu and Postdoctoral Fellow Eric Viollier have been involved in upgrading the STEADYSED1 code and running sensitivity simulations. Meanwhile Dr. Viollier's appointment at Georgia Tech has ended. In anticipation of continued funding for Phase II of the project, a new Postdoctoral fellow has been recruited to work on the stochastic bio-irrigation model: Dr. Carla Koretsky (Ph. D. 1997, Johns Hopkins University) will join the P.I.'s research group on January 1, 1998.

REFERENCES CITED IN REPORT

- Arakaki, T. and Mucci, A. (1995) A continuous and mechanistic representation of calcite reaction-controlled kinetics in dilute solution at 25°C and 1 atm total pressure. *Aquatic Geochem.*, 1, 1-26.
- Cai W.-J., Reimers C. E., and Shaw T. (1995) Microelectrode studies of organic carbon degradation and calcite dissolution at a California continental rise site. *Geochim. Cosmochim. Acta*, 59, 497-511.
- Canfield D. E., Thamdrup B., and Hansen J. W. (1993a) The anaerobic degradation of organic matter in Danish coastal sediments: Iron reduction, manganese reduction, and sulfate reduction. *Geochim. Cosmochim. Acta*, 57, 3867-3883.
- Canfield D. E., Jørgensen B. B., Fossing H., Glud R., Gundersen J., Ramsing N. B., Thamdrup B., Hansen J. W., Nielsen L. P., and Hall P. O. J. (1993b) Pathways of organic carbon oxidation in three continental margin sediments. *Mar. Geology*, 113, 27-40.
- Gaillard, J.-F., Pauwels, H., and Michard, G. (1989) Chemical diagenesis in coastal marine sediments. *Oceanol. Acta*, 12, 175-187.
- Chou, L., Garrels, R. M., and Wollast, R. (1989) Comparative study of the kinetics and mechanisms of dissolution of carbonate minerals. *Chem. Geology*, 78, 269-282.

- Martin W. R. and Banta G. T. (1992) The measurement of sediment irrigation rates: A comparison of the Br⁻ tracer and ²²²Rn/²²⁶Ra disequilibrium techniques. *J. Marine Res.*, 50, 125-154.
- Middelburg, J. J., Soetaert, K. and Herman, P. M. (1007) Empirical relationships for use in global diagenetic models. *Deep-Sea Res. I*, 44, 327-344.
- Plummer, L. N., Wigley, T. M. L., and Parkhurst, D. L. (1979) Critical review of the kinetics of calcite dissolution and precipitation. In *Chemical Modeling of Aqueous Systems: Speciation, Sorption, Solubility, and Kinetics* (Ed. E. A. Jenne). Amer. Chem. Soc. Symp. Ser., 93, 537-573.
- Tromp T. K., Van Cappellen P., and Key R. M. (1995) A global model for the early diagenesis of organic carbon and organic phosphorus in marine sediments. *Geochim. Cosmochim. Acta*, 59, 1259-1284.
- Van Cappellen P., Gaillard J.-F., and Rabouille C. (1993) Biogeochemical transformations in sediments: Kinetic models of early diagenesis. In *Interactions of C, N, P, and S Biogeochemical Cycles and Global Change* (ed. R. Wollast, F. T. Mackenzie, and L. Chou), pp. 401-445. Springer-Verlag.
- Wang, Y. and Van Cappellen, P. (1996) A multi-component reaction-transport model of early diagenesis: Application to redox cycling in coastal marine sediments. *Geochim. Cosmochim. Acta*, 60, 2993-3014.